

<p>Drug Facts</p>	<p>Drug Facts (continued)</p>
<p>Active ingredient (in each tablet) Purpose Aspirin 81 mg (NSAID)*Pain reliever *nonsteroidal anti-inflammatory drug</p>	<p>Do not use</p> <ul style="list-style-type: none"> ■ if you are allergic to aspirin or any other pain reliever/fever reducer ■ for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor
<p>Uses</p> <ul style="list-style-type: none"> ■ temporarily relieves minor aches and pains ■ ask your doctor about other uses for St. Joseph 81 mg Aspirin 	<p>Ask a doctor before use if</p> <ul style="list-style-type: none"> ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma
<p>Warnings</p> <p>Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.</p> <p>Allergy alert: Aspirin may cause a severe allergic reaction which may include:</p> <ul style="list-style-type: none"> ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock <p>Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you</p> <ul style="list-style-type: none"> ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed 	<p>Ask a doctor or pharmacist before use if you are taking a prescription drug for gout, diabetes or arthritis</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> ■ an allergic reaction occurs. Seek medical help right away. ■ you experience any of the following signs of stomach bleeding: <ul style="list-style-type: none"> ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ pain gets worse or lasts for more than 10 days ■ ringing in the ears or loss of hearing occurs ■ redness or swelling is present ■ new symptoms occur. These could be signs of a serious condition. <p>If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.</p>

<p>Drug Facts (continued)</p>	<p>Drug Facts (continued)</p>
<p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p>	<p>Other information</p> <ul style="list-style-type: none"> ■ store at 20-25°C (68-77°F) ■ close cap tightly after use
<p>Directions</p> <ul style="list-style-type: none"> ■ drink a full glass of water with each dose ■ adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor ■ children under 12 years: do not use unless directed by a doctor 	<p>Inactive ingredients</p> <p>corn starch, dextrates hydrated, FD&C yellow #6 aluminum lake, flavor, maltodextrin, sucralose</p>
	<p>Questions or comments?</p> <p>Call 1-855-785-6381 weekdays 9 AM to 5 PM EST</p>