

## Drug Facts

### Active ingredient

#### (in each tablet) Purpose

Aspirin 81 mg (NSAID)\* .....Pain  
reliever

\*nonsteroidal anti-inflammatory drug

### Uses

- temporarily relieves minor aches and pains
- ask your doctor about other uses for St. Joseph 81 mg Aspirin

### Warnings

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

## Drug Facts (continued)

### Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor

### Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

### Ask a doctor or pharmacist before use if you are taking a prescription drug for gout, diabetes or arthritis

### Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- pain gets worse or lasts for more than 10 days
- ringing in the ears or loss of hearing occurs
- redness or swelling is present
- new symptoms occur. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

## Drug Facts (continued)

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

### Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

## Drug Facts (continued)

### Other information

- store at 20-25°C (68-77°F)
- close cap tightly after use

### Inactive ingredients

corn starch, dextrans hydrated, FD&C yellow #6 aluminum lake, flavor, maltodextrin, sucralose

### Questions or comments?

Call 1-855-785-6381  
weekdays 9 AM to 5 PM EST